

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85934 A1

(51) International Patent Classification⁷: C12N 15/12,
15/62, C07K 14/47, 16/18, C12Q 1/68, G01N 33/68

Christian [DE/DE]; Heinrich-Delp-Strasse 221, 64297
Darmstadt (DE).

(21) International Application Number: PCT/EP01/04950

(74) Common Representative: MERCK PATENT GMBH;
Frankfurter Strasse 250, 64293 Darmstadt (DE).

(22) International Filing Date: 3 May 2001 (03.05.2001)

(81) Designated States (*national*): CA, JP, US.

(25) Filing Language: English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(26) Publication Language: English

(30) Priority Data:
00109133.9 5 May 2000 (05.05.2000) EP

Published:
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant (*for all designated States except US*): MERCK
PATENT GMBH [DE/DE]; Frankfurter Strasse 250,
64293 Darmstadt (DE).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): SIRRENBURG,



WO 01/85934 A1

(54) Title: NEW RAS GUANINE-NUCLEOTIDE-EXCHANGE FACTOR 1 (NRG1)

(57) Abstract: New Ras Guanine-nucleotide-exchange factor 1 (NRG1) polypeptides and polynucleotides and methods for produc-
ing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing New Ras Guanine-nu-
cleotide-exchange factor 1 (NRG1) polypeptides and polynucleotides in diagnostic assays.

New Ras Guanine-nucleotide-exchange factor 1 (NRG1)

Field of the Invention

5 This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides sometimes hereinafter referred to as „Ras Guanine-nucleotide-exchange factor 1 (NRG1)“, to their use in diagnosis and in identifying compounds that may be agonists, antagonists that are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

Background of the Invention

10 The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superseding earlier approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

15 Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Summary of the Invention

20 The present invention relates to New Ras Guanine-nucleotide-exchange factor 1 (NRG1), in particular New Ras Guanine-nucleotide-exchange factor 1 (NRG1)polypeptides and New Ras Guanine-nucleotide-exchange factor 1 (NRG1)polynucleotides, recombinant materials and methods for

25

30

their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain diseases, including, but not limited to, cancer, hereinafter referred to as "diseases of the invention". In a further aspect, the invention relates to methods for identifying agonists and antagonists (e.g., inhibitors) using the materials provided by the invention, and treating conditions associated with New Ras Guanine-nucleotide-exchange factor 1 (NRG1) imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate New Ras Guanine-nucleotide-exchange factor 1 (NRG1) activity or levels.

Description of the Invention

In a first aspect, the present invention relates to New Ras Guanine-nucleotide-exchange factor 1 (NRG1) polypeptides. Such polypeptides include:

(a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 3;

(b) a polypeptide comprising a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;

(c) a polypeptide comprising the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;

(d) a polypeptide having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;

(e) the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4; and

(f) a polypeptide having or comprising a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;

(g) fragments and variants of such polypeptides in (a) to (f).

Polypeptides of the present invention are believed to be members of the Ras guanine-nucleotide-exchange factor (GEFs) family of polypeptides.

They are therefore of interest because Ras guanine-nucleotide-exchange factors regulate the activation of Ras-like proteins (Boguski, M.S. and McCormick, F. (1993), Nature 366, 643-654). Ras proteins and their relatives play critical roles in the control of cell growth and differentiation and morphology. Ras is a signal-transducing, guanine nucleotide-binding protein for various membrane receptors including tyrosine kinase receptors and G-protein-coupled receptors (Rozakis-Adcock-M; et al. (1993), Nature 363, 83-85; Insall, R.H. et al. (1996), Curr. Biol. 6, 719-729, Kiyono, M. et al. (1999), PNAS 96, 4826-4831). Ras-like proteins function as molecular switches that cycle between GTP-bound ON and GDP bound OFF state. Binding of GTP (ON state) allows the GTP-binding proteins to interact with effector molecules. Hydrolysis of bound GTP is the timing mechanism that returns these proteins to their GDP-bound (OFF) state. In this way GEFs turn on the switches (activate Ras-like proteins) by exchanging bound GDP for free GTP. The importance of GTPase regulation is evident from diseases associated with mutations in the oncogene ras, which lead to constitutive activation of the Ras protein and contribute to malignant transformation.

The biological properties of the New Ras Guanine-nucleotide-exchange factor 1 (NRG1) are hereinafter referred to as "biological activity of New Ras Guanine-nucleotide-exchange factor 1 (NRG1)" or "New Ras Guanine-nucleotide-exchange factor 1 (NRG1) activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of New Ras Guanine-nucleotide-exchange factor 1 (NRG1).

Polypeptides of the present invention also includes variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

Preferred fragments of polypeptides of the present invention include a polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4, or a polypeptide comprising an amino acid

sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4. Preferred fragments are biologically active fragments that mediate the biological activity of New Ras Guanine-nucleotide-exchange factor 1 (NRG1), including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods. Means for preparing such polypeptides are well understood in the art.

In a further aspect, the present invention relates to New Ras Guanine-nucleotide-exchange factor 1 (NRG1) polynucleotides. Such polynucleotides include:

(a) a polynucleotide comprising a polynucleotide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3;

(b) a polynucleotide comprising the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;

(c) a polynucleotide having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;

- (d) the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;
- (e) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;
- 5 (f) a polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4;
- (g) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;
- 10 (h) a polynucleotide encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4;
- (i) a polynucleotide having or comprising a polynucleotide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3;
- 15 (j) a polynucleotide having or comprising a polynucleotide sequence encoding a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4; and
- 20 polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.
- Preferred fragments of polynucleotides of the present invention include a polynucleotide comprising a nucleotide sequence having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1 or
- 25 SEQ ID NO: 3, or a polynucleotide comprising a sequence having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ ID NO: 1 or SEQ ID NO: 3.
- Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including
- 30 polynucleotides having one or more single nucleotide polymorphisms (SNPs).

Polynucleotides of the present invention also include polynucleotides encoding polypeptide variants that comprise the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 4 and in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acid residues are substituted, deleted or added, in any combination.

In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide that:

- (a) comprises an RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4;
- (b) is the RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4;
- (c) comprises an RNA transcript of the DNA sequence of SEQ ID NO: 1 or SEQ ID NO: 3; or
- (d) is the RNA transcript of the DNA sequence of SEQ ID NO: 1 or SEQ ID NO: 3;

and RNA polynucleotides that are complementary thereto.

The polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3 shows homology with CG7369 (*D. melongaster*, Acc. AE003598), Ro5G6.10 (*C. elegans*, Acc. U58746), members of the RasGEF protein family including the aimless RasGEF (*Dictyostelium discoideum*, Acc. U53884), "sterile" gene 6 (Ste6, *S. pombe*, Acc. X53254), SCD25 (SC25, *S. cerevisiae*, Acc. X03579), son of sevenless (SOS, *D. melongaster*, Acc. M83931), guanine nucleotide dissociation stimulator A (GNDS, mouse, Acc. L07924). The polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3 is a cDNA sequence that encodes the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4. The polynucleotide sequence encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4 may be identical to the polypeptide encoding sequence of SEQ ID NO: 1 or SEQ ID NO: 3 or it may be a sequence other than SEQ ID NO: 1 or SEQ ID NO: 3, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4. The polypeptide of the SEQ ID NO: 2 or SEQ

ID NO: 4 is related to other proteins of the Ras guanine-nucleotide-exchange factor (GEFs) family, having homology and/or structural similarity with members of the RasGEF protein family.

Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one New Ras Guanine-nucleotide-exchange factor 1 (NRG1) activity.

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human colon carcinoma (see for instance, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and

primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO: 1 or SEQ ID NO: 3, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or SEQ ID NO: 3 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes isolated polynucleotides, preferably with a nucleotide sequence of at least 100, obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or SEQ ID NO: 3 or a fragment thereof, preferably of at least 15 nucleotides.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., Proc Nat Acad Sci USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trade mark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark) technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adapter specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook *et al.* (*ibid*). Preferred methods of introducing polynucleotides into host cells include, for

instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, micro-injection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

5 Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells;
10 and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements,
15 from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control
20 regions that regulate as well as engender expression. Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as,
25 for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

30 If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to
35 recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or purification.

Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterized by the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled New Ras Guanine-nucleotide-exchange factor 1 (NRG1)nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the

chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401).

5 An array of oligonucleotides probes comprising New Ras Guanine-nucleotide-exchange factor 1 (NRG1) polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability, see, for
10 example, M. Chee *et al.*, Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or
15 increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a
20 protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radio-immunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

25 Thus in another aspect, the present invention relates to a diagnostic kit comprising:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3, or a fragment or an RNA transcript thereof;
- (b) a nucleotide sequence complementary to that of (a);
- 30 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

5 The polynucleotide sequences of the present invention are valuable for chromosome localisation studies. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating
10 those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical
15 Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (co-inheritance of physically adjacent genes). Precise human chromosomal localisations for a genomic sequence (gene fragment etc.) can be determined using Radiation Hybrid (RH) Mapping (Walter, M. Spillett, D., Thomas, P., Weissenbach, J., and Goodfellow, P.,
20 (1994) A method for constructing radiation hybrid maps of whole genomes, Nature Genetics 7, 22-28). A number of RH panels are available from Research Genetics (Huntsville, AL, USA) e.g. the GeneBridge4 RH panel (Hum Mol Genet 1996 Mar;5(3):339-46 A radiation hybrid map of the human genome. Gyapay G, Schmitt K, Fizames C, Jones H, Vega-Czarny N, Spillett D, Muselet D, Prud'Homme JF, Dib C, Auffray C, Morissette J, Weissenbach J, Goodfellow PN). To determine the chromosomal location of a gene using this panel, 93 PCRs are performed using primers designed from the gene of interest on RH
25 DNAs. Each of these DNAs contains random human genomic fragments maintained in a hamster background (human / hamster hybrid cell lines). These PCRs result in 93 scores indicating the presence or absence of the PCR product of the gene of interest. These scores are compared with scores created using PCR products from genomic sequences of known location. This comparison is conducted at
35 <http://www.genome.wi.mit.edu/>.

The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridization techniques to clones arrayed on a grid, such as cDNA microarray hybridization (Schena *et al*, Science, 270, 467-470, 1995 and Shalon *et al*, Genome Res, 6, 639-645, 1996) and nucleotide amplification techniques such as PCR. A preferred method uses the TAQMAN (Trade mark) technology available from Perkin Elmer. Results from these studies can provide an indication of the normal function of the polypeptide in the organism. In addition, comparative studies of the normal expression pattern of mRNAs with that of mRNAs encoded by an alternative form of the same gene (for example, one having an alteration in polypeptide coding potential or a regulatory mutation) can provide valuable insights into the role of the polypeptides of the present invention, or that of inappropriate expression thereof in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

The polypeptides of the present invention are expressed in colon carcinoma..

A further aspect of the present invention relates to antibodies. The polypeptides of the invention or their fragments, or cells expressing them, can be used as immunogens to produce antibodies that are immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., Nature (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, Immunology Today (1983) 4:72) and

the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease, whether that disease is already established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intra-muscular, intravenous, or intra-dermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile

suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include
5 adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

10 Polypeptides of the present invention have one or more biological functions that are of relevance in one or more disease states, in particular the diseases of the invention hereinbefore mentioned. It is therefore useful to identify compounds that stimulate or inhibit the function or level of the
15 polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those that stimulate or inhibit the function or level of the polypeptide. Such methods identify agonists or antagonists that may be employed for therapeutic and prophylactic purposes for such diseases of the invention as hereinbefore
20 mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, collections of chemical compounds, and natural product mixtures. Such agonists or antagonists so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; a
25 structural or functional mimetic thereof (see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)) or a small molecule. Such small molecules preferably have a molecular weight below 2,000 daltons, more preferably between 300 and 1,000 daltons, and most preferably between 400 and 700 daltons. It is preferred that these small molecules
30 are organic molecules.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof, by means of a label directly or indirectly associated with the candidate compound. Alternatively, the
35 screening method may involve measuring or detecting (qualitatively or quantitatively) the competitive binding of a candidate compound to the

polypeptide against a labeled competitor (e.g. agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring a New Ras Guanine-nucleotide-exchange factor 1 (NRG1) activity in the mixture, and comparing the New Ras Guanine-nucleotide-exchange factor 1 (NRG1) activity of the mixture to a control mixture which contains no candidate compound.

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well micotiter plates but also emerging methods such as the nanowell method described by Schullek et al, Anal Biochem., 246, 20-29, (1997).

Fusion proteins, such as those made from Fc portion and New Ras Guanine-nucleotide-exchange factor 1 (NRG1) polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

Screening techniques

The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents that may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

A polypeptide of the present invention may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide that compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

Examples of antagonists of polypeptides of the present invention include antibodies or, in some cases, oligonucleotides or proteins that are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or a small molecule that bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Screening methods may also involve the use of transgenic technology and New Ras Guanine-nucleotide-exchange factor 1 (NRG1) gene. The art of constructing transgenic animals is well established. For example, the New Ras Guanine-nucleotide-exchange factor 1 (NRG1) gene may be introduced through microinjection into the male pronucleus of fertilized oocytes, retroviral transfer into pre- or post-implantation embryos, or injection of genetically modified, such as by electroporation, embryonic stem cells into host blastocysts. Particularly useful transgenic animals are so-called "knock-in" animals in which an animal gene is replaced by the human equivalent within the genome of that animal. Knock-in transgenic animals are useful in the drug discovery process, for target validation, where the compound is specific for the human target. Other useful transgenic animals are so-called "knock-out" animals in which the expression of the animal ortholog of a polypeptide of the present invention and encoded by an endogenous DNA sequence in a cell is partially or completely annulled. The gene knock-out may be targeted to specific cells or tissues, may occur only in certain cells or tissues as a

consequence of the limitations of the technology, or may occur in all, or substantially all, cells in the animal. Transgenic animal technology also offers a whole animal expression-cloning system in which introduced genes are expressed to give large amounts of polypeptides of the present invention

Screening kits for use in the above described methods form a further aspect of the present invention. Such screening kits comprise:

- (a) a polypeptide of the present invention;
- (b) a recombinant cell expressing a polypeptide of the present invention;
- (c) a cell membrane expressing a polypeptide of the present invention; or
- (d) an antibody to a polypeptide of the present invention;

which polypeptide is preferably that of SEQ ID NO: 2 or SEQ ID NO: 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an

Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by

transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

5 "Polynucleotide" generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules
10 comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and
15 DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in
20 nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds,
25 i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural
30 processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide
35 backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide.

Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, 1-12, in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol*, 182, 626-646, 1990, and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci*, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO: 1 or SEQ ID NO: 3.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions,

deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more)

PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript undergoes splicing, generally for the removal of introns, which results in the production of more than one mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

"Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over the length of the sequences being compared.

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two polypeptide chains, on a residue by residue basis, taking into account not only exact correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an

associated "score" from which the "% similarity" of the two sequences can then be determined.

Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J et al, Nucleic Acids Res, 12, 387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the % similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (J Mol Biol, 147,195-197, 1981, Advances in Applied Mathematics, 2, 482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Needleman and Wunsch (J Mol Biol, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F et al, J Mol Biol, 215, 403-410, 1990, Altschul S F et al, Nucleic Acids Res., 25:389-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at www.ncbi.nlm.nih.gov) and FASTA (Pearson W R, Methods in Enzymology, 183, 63-99, 1990; Pearson W R and Lipman D J, Proc Nat Acad Sci USA, 85, 2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package).

Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, Proc. Nat. Acad Sci. USA, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

"Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid

5 deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between these terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

15 The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \bullet I),$$

in which:

n_a is the number of nucleotide or amino acid differences,

20 x_a is the total number of nucleotides or amino acids in SEQ ID NO: 1 or SEQ ID NO: 3 or SEQ ID NO: 2 or SEQ ID NO: 4, respectively,

I is the Identity Index,

\bullet is the symbol for the multiplication operator, and

in which any non-integer product of x_a and I is rounded down to the nearest integer prior to subtracting it from x_a .

25 "Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the two sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or

30

polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

5 "Fusion protein" refers to a protein encoded by two, unrelated, fused genes or fragments thereof. Examples have been disclosed in US 5541087, 5726044. In the case of Fc-NRG1, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for performing the functional expression of Fc-NRG1 or fragments of NRG1, to improve pharmacokinetic properties of such a fusion protein when used for therapy and to generate a dimeric NRG1. The Fc- NRG1
10 DNA construct comprises in 5' to 3' direction, a secretion cassette, i.e. a signal sequence that triggers export from a mammalian cell, DNA encoding an immunoglobulin Fc region fragment, as a fusion partner, and a DNA encoding NRG1 or fragments thereof. In some uses it would be desirable to be able to alter the intrinsic functional properties
15 (complement binding, Fc-Receptor binding) by mutating the functional Fc sides while leaving the rest of the fusion protein untouched or delete the Fc part completely after expression.

20 All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and
25 references.

Claims

1. A polypeptide selected from the group consisting of:
 - (a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 3;
 - (b) a polypeptide comprising a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;
 - (c) a polypeptide having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;
 - (d) the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4 and
 - (e) fragments and variants of such polypeptides in (a) to (d).
2. The polypeptide of claim 1 comprising the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4.
3. The polypeptide of claim 1 which is the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4.
4. A polynucleotide selected from the group consisting of:
 - (a) a polynucleotide comprising a polynucleotide sequence having at least 95% identity to the polynucleotide sequence of SEQ ID NO: 1 or SEQ ID NO: 3;
 - (b) a polynucleotide having at least 95% identity to the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;
 - (c) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;

- (d) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO: 2 or SEQ ID NO: 4;
- (e) a polynucleotide with a nucleotide sequence of at least 100 nucleotides
5 obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or SEQ ID NO: 3 or a fragment thereof having at least 15 nucleotides;
- (f) a polynucleotide which is the RNA equivalent of a polynucleotide of (a) to (e);
- (g) a polynucleotide sequence complementary to said polynucleotide of any one of
10 (a) to (f), and
- (h) polynucleotides that are variants or fragments of the polynucleotides of any one of (a) to (g) or that are complementary to above mentioned polynucleotides, over the entire length thereof.

15 5. A polynucleotide of claim 4 selected from the group consisting of:

- (a) a polynucleotide comprising the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;
- (b) the polynucleotide of SEQ ID NO: 1 or SEQ ID NO: 3;
- (c) a polynucleotide comprising a polynucleotide sequence encoding the
20 polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4; and
- (d) a polynucleotide encoding the polypeptide of SEQ ID NO: 2 or SEQ ID NO: 4.

6. An expression system comprising a polynucleotide capable of producing a polypeptide of any one of claim 1-3 when said expression vector is present in a
25 compatible host cell.

7. A recombinant host cell comprising the expression vector of claim 6 or a membrane thereof expressing the polypeptide of any one of claim 1-3.

8. A process for producing a polypeptide of any one of claim 1-3 comprising the step of culturing a host cell as defined in claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
9. A fusion protein consisting of the Immunoglobulin Fc-region and a polypeptide any one one of claims 1-3.
10. An antibody immunospecific for the polypeptide of any one of claims 1 to 3.
11. A method for screening to identify compounds that stimulate or inhibit the function or level of the polypeptide of any one of claim 1-3 comprising a method selected from the group consisting of:
- (a) measuring or, detecting, quantitatively or qualitatively, the binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;
 - (b) measuring the competition of binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
 - (c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes expressing the polypeptide;
 - (d) mixing a candidate compound with a solution containing a polypeptide of any one of claims 1-3, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a control mixture which contains no candidate compound; or

(e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide or said polypeptide in cells, using for instance, an ELISA assay, and

(f) producing said compound according to biotechnological or chemical standard techniques.

SEQUENCE LISTING

<110> Merck Patent GmbH

5 <120> Ras Guanine-nucleotide -exchange factor (NRG1)

<130> NRG1CSWS

<140>

10 <141>

<160> 4

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 1422

<212> DNA

<213> Homo sapiens

20

<220>

<221> CDS

<222> (1)..(1422)

25 <400> 1

atg cct cag act cct ccc ttt tca gca atg ttt gac agc agt ggt tac	48
Met Pro Gln Thr Pro Pro Phe Ser Ala Met Phe Asp Ser Ser Gly Tyr	
1 5 10 15	

aat cga aac ctc tat cag tct gca gag gac agc tgt gga ggg ttg tat	96
Asn Arg Asn Leu Tyr Gln Ser Ala Glu Asp Ser Cys Gly Gly Leu Tyr	
20 25 30	

tac cat gac aac aac ctc ctc tct gga tcc ctg gaa gca ctc atc cag	144
Tyr His Asp Asn Asn Leu Leu Ser Gly Ser Leu Glu Ala Leu Ile Gln	
35 40 45	

cac tta gta cct aat gtg gat tac tat cca gat aga aca tac ata ttt	192
His Leu Val Pro Asn Val Asp Tyr Tyr Pro Asp Arg Thr Tyr Ile Phe	
40 50 55 60	

acc ttc cta ctc agt tct cgg tta ttt atg cat ccg tat gag cta atg	240
Thr Phe Leu Leu Ser Ser Arg Leu Phe Met His Pro Tyr Glu Leu Met	
65 70 75 80	

gcc aaa gtt tgc cac tta tgt gtt gag cac cag aga cta agt gat cct	288
Ala Lys Val Cys His Leu Cys Val Glu His Gln Arg Leu Ser Asp Pro	
85 90 95	

gat agt gat aag aac cag atg aga aaa att gca ccc aaa atc ctt caa	336
Asp Ser Asp Lys Asn Gln Met Arg Lys Ile Ala Pro Lys Ile Leu Gln	
100 105 110	

ctc ctc acg gaa tgg acg gaa aca ttt ccc tat gat ttt cgg gat gaa	384
Leu Leu Thr Glu Trp Thr Glu Thr Phe Pro Tyr Asp Phe Arg Asp Glu	
115 120 125	

aga atg atg aga aac tta aaa gat ctg gct cac cga ata gcc agt ggc	432
Arg Met Met Arg Asn Leu Lys Asp Leu Ala His Arg Ile Ala Ser Gly	
60 130 135 140	

	gaa gag cag aca tac aga aag aat gtc cag caa atg atg cag tgt ctg	480
	Glu Glu Gln Thr Tyr Arg Lys Asn Val Gln Gln Met Met Gln Cys Leu	
	145 150 155 160	
5	atc cgc aag ctt gct gcg ctc agc cag tac gaa gaa gtc ctg gca aaa	528
	Ile Arg Lys Leu Ala Ala Leu Ser Gln Tyr Glu Glu Val Leu Ala Lys	
	165 170 175	
10	atc agc tcc aca tcc aca gat cgg ctc aca gtt ctc aag acc aag cca	576
	Ile Ser Ser Thr Ser Thr Asp Arg Leu Thr Val Leu Lys Thr Lys Pro	
	180 185 190	
15	cag tct ata caa agg gat atc att act gtc tgc aac gac cct tac acg	624
	Gln Ser Ile Gln Arg Asp Ile Ile Thr Val Cys Asn Asp Pro Tyr Thr	
	195 200 205	
20	ttg gcc cag cag ctg act cat ata gag ctg gag agg ctc aat tat att	672
	Leu Ala Gln Gln Leu Thr His Ile Glu Leu Glu Arg Leu Asn Tyr Ile	
	210 215 220	
25	ggg cca gaa gaa ttt gtt cag gcg ttc gtg cag aag gac cct ttg gat	720
	Gly Pro Glu Glu Phe Val Gln Ala Phe Val Gln Lys Asp Pro Leu Asp	
	225 230 235 240	
	aat gac aag agt tgc tac agt gaa cgg aag aaa aca cga aac tta gaa	768
	Asn Asp Lys Ser Cys Tyr Ser Glu Arg Lys Lys Thr Arg Asn Leu Glu	
	245 250 255	
30	gct tac gtg gaa tgg ttt aat cgc ctc agc tac ttg gtt gct aca gaa	816
	Ala Tyr Val Glu Trp Phe Asn Arg Leu Ser Tyr Leu Val Ala Thr Glu	
	260 265 270	
35	atc tgt atg cct gtt aag aaa aaa cac cga gca aga atg att gag tat	864
	Ile Cys Met Pro Val Lys Lys Lys His Arg Ala Arg Met Ile Glu Tyr	
	275 280 285	
40	ttc att gac gta gct cgg gag tgt ttt aac att ggc aac ttc aac tcc	912
	Phe Ile Asp Val Ala Arg Glu Cys Phe Asn Ile Gly Asn Phe Asn Ser	
	290 295 300	
45	ttg atg gcg ata atc tct ggt atg aat atg agc cca gtc tct cga cta	960
	Leu Met Ala Ile Ile Ser Gly Met Asn Met Ser Pro Val Ser Arg Leu	
	305 310 315 320	
	aaa aaa act tgg gcc aaa gtg aag act gca aaa ttt gac att ctt gag	1008
	Lys Lys Thr Trp Ala Lys Val Lys Thr Ala Lys Phe Asp Ile Leu Glu	
	325 330 335	
50	cat cag atg gac cct tca agc aat ttc tat aat tat cga aca gct ctt	1056
	His Gln Met Asp Pro Ser Ser Asn Phe Tyr Asn Tyr Arg Thr Ala Leu	
	340 345 350	
55	cgt ggg gca gca caa agg tct tta act gct cat agt agt aga gaa aag	1104
	Arg Gly Ala Ala Gln Arg Ser Leu Thr Ala His Ser Ser Arg Glu Lys	
	355 360 365	
60	att gtg ata cca ttc ttc agt ctc tta atc aaa gat att tat ttc ctc	1152
	Ile Val Ile Pro Phe Phe Ser Leu Leu Ile Lys Asp Ile Tyr Phe Leu	
	370 375 380	

aat gag ggt tgt gcc aac cgc ctt ccc aat ggc cat gtc aat ttt gag 1200
 Asn Glu Gly Cys Ala Asn Arg Leu Pro Asn Gly His Val Asn Phe Glu
 385 390 395 400

5 aaa ttt tgg gaa ctg gcc aaa caa gtg agt gaa ttt atg aca tgg aaa 1248
 Lys Phe Trp Glu Leu Ala Lys Gln Val Ser Glu Phe Met Thr Trp Lys
 405 410 415

10 caa gtg gag tgt cca ttt gag agg gac cgg aag atc ttg cag tat ctg 1296
 Gln Val Glu Cys Pro Phe Glu Arg Asp Arg Lys Ile Leu Gln Tyr Leu
 420 425 430

15 ctc aca gta cca gtc ttc agt gaa gat gct ctc tac ttg gct tct tat 1344
 Leu Thr Val Pro Val Phe Ser Glu Asp Ala Leu Tyr Leu Ala Ser Tyr
 435 440 445

gag agt gaa gga cct gaa aat cat ata gag aaa gac aga tgg aag tct 1392
 Glu Ser Glu Gly Pro Glu Asn His Ile Glu Lys Asp Arg Trp Lys Ser
 450 455 460

20 tta agg tcg agc ctc tta ggc aga gtt taa 1422
 Leu Arg Ser Ser Leu Leu Gly Arg Val
 465 470

25
 <210> 2
 <211> 473
 <212> PRT
 <213> Homo sapiens

30
 <400> 2
 Met Pro Gln Thr Pro Pro Phe Ser Ala Met Phe Asp Ser Ser Gly Tyr
 1 5 10 15
 Asn Arg Asn Leu Tyr Gln Ser Ala Glu Asp Ser Cys Gly Gly Leu Tyr
 20 25 30
 Tyr His Asp Asn Asn Leu Leu Ser Gly Ser Leu Glu Ala Leu Ile Gln
 35 40 45
 His Leu Val Pro Asn Val Asp Tyr Tyr Pro Asp Arg Thr Tyr Ile Phe
 50 55 60
 40 Thr Phe Leu Leu Ser Ser Arg Leu Phe Met His Pro Tyr Glu Leu Met
 65 70 75 80
 Ala Lys Val Cys His Leu Cys Val Glu His Gln Arg Leu Ser Asp Pro
 85 90 95
 Asp Ser Asp Lys Asn Gln Met Arg Lys Ile Ala Pro Lys Ile Leu Gln
 100 105 110
 45 Leu Leu Thr Glu Trp Thr Glu Thr Phe Pro Tyr Asp Phe Arg Asp Glu
 115 120 125
 Arg Met Met Arg Asn Leu Lys Asp Leu Ala His Arg Ile Ala Ser Gly
 130 135 140
 50 Glu Glu Gln Thr Tyr Arg Lys Asn Val Gln Gln Met Met Gln Cys Leu
 145 150 155 160
 Ile Arg Lys Leu Ala Leu Ser Gln Tyr Glu Glu Val Leu Ala Lys
 165 170 175
 Ile Ser Ser Thr Ser Thr Asp Arg Leu Thr Val Leu Lys Thr Lys Pro
 180 185 190
 55 Gln Ser Ile Gln Arg Asp Ile Ile Thr Val Cys Asn Asp Pro Tyr Thr
 195 200 205
 Leu Ala Gln Gln Leu Thr His Ile Glu Leu Glu Arg Leu Asn Tyr Ile
 210 215 220
 60 Gly Pro Glu Glu Phe Val Gln Ala Phe Val Gln Lys Asp Pro Leu Asp
 225 230 235 240

[illegible]

```

35  <210> 3
    <211> 423
    <212> DNA
    <213> Homo sapiens

```

```

    <220>
40    <221> CDS
    <222> (1) .. (423)

```

	<400> 3	
	caa agg gat atc att act gtc tgc aac gac cct tac acg ttg gcc cag	48
45	Gln Arg Asp Ile Ile Thr Val Cys Asn Asp Pro Tyr Thr Leu Ala Gln	
	1 5 10 15	
	cag ctg act cat ata gag ctg gag agg ctc aat tat att ggg cca gaa	96
	Gln Leu Thr His Ile Glu Leu Glu Arg Leu Asn Tyr Ile Gly Pro Glu	
50	20 25 30	
	gaa ttt gtt cag gcg ttc gtg cag aag gac cct ttg gat aat gac aag	144
	Glu Phe Val Gln Ala Phe Val Gln Lys Asp Pro Leu Asp Asn Asp Lys	
	35 40 45	
55	agt tgc tac agt gaa cgg aag aaa aca cga aac tta gaa gct tac gtg	192
	Ser Cys Tyr Ser Glu Arg Lys Lys Thr Arg Asn Leu Glu Ala Tyr Val	
	50 55 60	

gaa tgg ttt aat cgc ctc agc tac ttg gtt gct aca gaa atc tgt atg 240
 Glu Trp Phe Asn Arg Leu Ser Tyr Leu Val Ala Thr Glu Ile Cys Met
 65 70 75 80

5 cct gtt aag aaa aaa cac cga gca aga atg att gag tat ttc att gac 288
 Pro Val Lys Lys Lys His Arg Ala Arg Met Ile Glu Tyr Phe Ile Asp
 85 90 95

gta gct cgg gag tgt ttt aac att ggc aac ttc aac tcc ttg atg gcg 336
 10 Val Ala Arg Glu Cys Phe Asn Ile Gly Asn Phe Asn Ser Leu Met Ala
 100 105 110

ata atc tct ggt atg aat atg agc cca gtc tct cga cta aaa aaa act 384
 15 Ile Ile Ser Gly Met Asn Met Ser Pro Val Ser Arg Leu Lys Lys Thr
 115 120 125

tgg gcc aaa gtg aag act gca aaa ttt gac att ctt gag 423
 Trp Ala Lys Val Lys Thr Ala Lys Phe Asp Ile Leu Glu
 130 135 140

20

<210> 4
 <211> 141
 <212> PRT
 25 <213> Homo sapiens

<400> 4
 Gln Arg Asp Ile Ile Thr Val Cys Asn Asp Pro Tyr Thr Leu Ala Gln
 1 5 10 15

30 Gln Leu Thr His Ile Glu Leu Glu Arg Leu Asn Tyr Ile Gly Pro Glu
 20 25 30

Glu Phe Val Gln Ala Phe Val Gln Lys Asp Pro Leu Asp Asn Asp Lys
 35 35 40 45

Ser Cys Tyr Ser Glu Arg Lys Lys Thr Arg Asn Leu Glu Ala Tyr Val
 50 55 60

40 Glu Trp Phe Asn Arg Leu Ser Tyr Leu Val Ala Thr Glu Ile Cys Met
 65 70 75 80

Pro Val Lys Lys Lys His Arg Ala Arg Met Ile Glu Tyr Phe Ile Asp
 85 90 95

45 Val Ala Arg Glu Cys Phe Asn Ile Gly Asn Phe Asn Ser Leu Met Ala
 100 105 110

Ile Ile Ser Gly Met Asn Met Ser Pro Val Ser Arg Leu Lys Lys Thr
 50 115 120 125

Trp Ala Lys Val Lys Thr Ala Lys Phe Asp Ile Leu Glu
 130 135 140

55

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/04950

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/62 C07K14/47 C07K16/18 C12Q1/68
G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EM_EST 'Online! EMBL; ID AW444812, AC AW444812, 20 February 2000 (2000-02-20) NATIONAL CANCER INSTITUTE, CANCER GENOME ANATOMY PROJECT (CGAP): "UI-H-BI3-ajy-d-11-0-UI.s1 NCI_CGAP_Sub5 Homo sapiens cDNA clone IMAGE:2733380 3', mRNA sequence" XP002178533 Note: 99.7% nt seq identity with SEQ ID NO:1 in 383 nt overlap (10-392:967-585), 99.2% aa seq identity of translated sequence with SEQ ID NO:2 in 127 aa overlap the whole document</p> <p style="text-align: center;">--- -/-</p>	1-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

26 September 2001

Date of mailing of the international search report

10/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

Intel. Application No

PCT/EP 01/04950

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EM_HTG 'Online! EMBL; ID AC016325, AC AC016325, 26 November 1999 (1999-11-26) BIRREN B ET AL.: "Homo sapiens clone RP11-29L8, WORKING DRAFT SEQUENCE, 9 unordered pieces" XP002178534 Note: 100.0% nt seq identity with SEQ ID NO:1 in 177 nt overlap (107043-107219:177-1), 100.0% aa seq identity of translated sequence with SEQ ID NO:2 in 59 aa overlap page 26</p>	4
A	<p>DATABASE EM_EST 'Online! EMBL; ID AI884701, AC AI884701, 28 July 1999 (1999-07-28) NATIONAL CANCER INSTITUTE / NATIONAL INSTITUTE OF NEUROLOGICAL DISORDERS AND STROKE, BRAIN TUMOR GENOME ANATOMY PROJECT : "w183e12.x1 NCI_CGAP_Brn25 Homo sapiens cDNA clone IMAGE:2431534 3' similar to TR:061605 061605 PUTATIVE GUANINE NUCLEOTIDE RELEASING FACTOR ;, mRNA sequence" XP002178535 Note: 65.2% nt seq identity with SEQ ID NO:1 in 313 nt overlap (334-646:1417-1105), 60.6% aa seq identity of translated sequence with SEQ ID NO:2 in 104 aa overlap the whole document</p>	1,4
A	<p>FAM N P ET AL.: "Cloning and characterization of Ras-GRF2, a novel guanine nucleotide exchange factor for Ras." MOLECULAR AND CELLULAR BIOLOGY, vol. 17, no. 3, 1997, pages 1396-1406, XP002178531 ISSN: 0270-7306 Note: 25.1% aa seq identity of Ras-GRF2 with SEQ ID NO:2 in 354 aa overlap (864-1184:104-451) abstract figure 1</p>	1,4
A	<p>US 5 656 595 A (TOCQUE BRUNO ET AL) 12 August 1997 (1997-08-12) Note: 26.4% aa seq identity of SEQ ID NO:4 with SEQ ID NO:2 in 367 aa overlap (155-484:97-451) the whole document</p>	1-11

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/04950

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	REBHUN J F ET AL.: "Identification and characterization of a new family of guanine nucleotide exchange factors for the Ras-related GTPase Ral." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 275, no. 18, 15 March 2000 (2000-03-15), pages 13406-13410, XP002178532 ISSN: 0021-9258 abstract	
A	BOGUSKI M S ET AL.: "Proteins regulating Ras and its relatives" NATURE, vol. 366, 16 December 1993 (1993-12-16), pages 643-654, XP002039500 ISSN: 0028-0836 cited in the application abstract page 645, left-hand column, line 4 -page 647, left-hand column, line 44 table 1	
P,X	WO 00 31263 A (INCYTE PHARMA INC ;AZIMZAI YALDA (US); YUE HENRY (US); BANDMAN OLG) 2 June 2000 (2000-06-02) Note: 100.0% nt seq identity of SEQ ID NO:40 with SEQ ID NO:1 in 744 nt overlap (1-744:679-1422), 100.0% aa seq identity of SEQ ID NO:11 with SEQ ID NO:2 in 199 aa overlap (1-199:275-473) page 5, line 16 -page 7, line 6 page 61, table 2: SEQ ID NO:11 claims 1-20	1,4,6-11
P,X	EP 1 033 401 A (GENSET SA) 6 September 2000 (2000-09-06) Note: 100.0% nt seq identity of SEQ ID NO:2093 with SEQ ID NO:1 in 234 nt overlap (145-378:1-234), 100.0% aa seq identity of SEQ ID NO: 6170 with SEQ ID NO:2 in 78 aa overlap (1-78:1-78) the whole document	1,4,6-11
P,X	WO 01 05970 A (INCYTE GENOMICS INC ;PATTERSON CHANDRA (US); AZIMZAI YALDA (US); R) 25 January 2001 (2001-01-25) Note: 73.7% nt seq identity of SEQ ID NO:132 with SEQ ID NO:1 in 840 nt overlap (568-1401:564-1400), 67.9% aa seq identity of SEQ ID NO:66 with SEQ ID NO:2 in 473 aa overlap (1-465:1-472) page 4, line 35 -page 9, line 27 page 93, table 2: SEQ ID NO:66 claims 1-28	1,4,6-11

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/04950

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EM_EST 'Online! EMBL; ID AW842798, AC AW842798, 23 May 2000 (2000-05-23) DIAS N E ET AL.: "MR2-CN0037-210200-101-b02 CN0037 Homo sapiens cDNA, mRNA sequence." XP002178536 Note: 100.0% nt seq identity with SEQ ID NO:1 in 507 nt overlap (1-507:1362-856), 100.0% aa seq identity of translated sequence with SEQ ID NO:2 in 169 aa overlap the whole document</p>	1,4
A	<p>-& PROCEEDINGS OF THE NATIONAL ACADEMY OF THE UNITED STATES, vol. 97, no. 7, 28 March 2000 (2000-03-28), pages 3491-3496, XP000996193 abstract page 3495, right-hand column, line 18-20</p>	
P,X	<p>DATABASE EM_EST 'Online! EMBL; ID BE503530, AC BE503530, 7 August 2000 (2000-08-07) NATIONAL CANCER INSTITUTE, CANCER GENOME ANATOMY PROJECT (CGAP): "hz87b03.x1 NCI_CGAP_Lu24 Homo sapiens cDNA clone IMAGE:3214925 3', mRNA sequence" XP002178537 Note: 100.0% nt seq identity with SEQ ID NO:1 in 205 nt overlap (96-300:1-205), 100.0% aa seq identity of translated sequence with SEQ ID NO:2 in 68 aa overlap the whole document</p>	1,4
T	<p>SPRANG S: "GEFs: master regulators of G-protein activation" TIBS TRENDS IN BIOCHEMICAL SCIENCES, vol. 26, no. 4, 1 April 2001 (2001-04-01), pages 266-267, XP004241856 ISSN: 0968-0004 the whole document</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int'l. Application No

PCT/EP 01/04950

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5656595	A	12-08-1997	FR 2690162 A1	22-10-1993
			CA 2131166 A1	28-10-1993
			EP 0637334 A1	08-02-1995
			WO 9321314 A1	28-10-1993
			JP 7505774 T	29-06-1995
WO 0031263	A	02-06-2000	AU 1744900 A	13-06-2000
			EP 1135484 A2	26-09-2001
			WO 0031263 A2	02-06-2000
EP 1033401	A	06-09-2000	EP 1033401 A2	06-09-2000
WO 0105970	A	25-01-2001	AU 6223700 A	05-02-2001
			WO 0105970 A2	25-01-2001